

STRONG CONVERGENCE FOR THE MODIFIED MANN'S ITERATION OF λ -STRICT PSEUDOCONTRACTION

Hongjun Wang¹ and Yisheng Song^{1,2}

1. College of Mathematics and Information Science, Henan Normal University,
XinXiang, P.R. China, 453007

2. Department of Applied Mathematics, The Hong Kong Polytechnic University,
Hung Hom, Kowloon, Hong Kong

Abstract. In this paper, for an λ -strict pseudocontraction T , we prove strong convergence of the modified Mann's iteration defined by

$$x_{n+1} = \beta_n u + \gamma_n x_n + (1 - \beta_n - \gamma_n)[\alpha_n T x_n + (1 - \alpha_n)x_n],$$

where $\{\alpha_n\}$, $\{\beta_n\}$ and $\{\gamma_n\}$ in $(0, 1)$ satisfy:

(i) $0 \leq \alpha_n \leq \frac{\lambda}{K^2}$ with $\liminf_{n \rightarrow \infty} \alpha_n(\lambda - K^2 \alpha_n) > 0$;

(ii) $\lim_{n \rightarrow \infty} \beta_n = 0$ and $\sum_{n=1}^{\infty} \beta_n = \infty$;

(iii) $\limsup_{n \rightarrow \infty} \gamma_n < 1$.

Our results unify and improve some existing results.

Key Words and Phrases: λ -strict pseudocontraction, modified Mann's iteration, 2-uniformly smooth Banach space.

2000 AMS Subject Classification: 49J40, 47H05, 47H04, 65J15, 47H10.

Throughout this paper, let E be a Banach space with the norm $\|\cdot\|$ and the dual space E^* and $\langle y, x^* \rangle$ denote the value of $x^* \in E^*$ at $y \in E$. The normalized duality mapping J from E into 2^{E^*} is defined by the following equation:

$$J(x) = \{x^* \in E^* : \langle x, x^* \rangle = \|x\| \|x^*\|, \|x\| = \|x^*\|\}.$$

Let $F(T) = \{x \in E : Tx = x\}$, the set of all fixed point of a mapping T .

Recall that a mapping T with domain $D(T)$ and range $R(T)$ in Banach space E is called *Lipschitzian* if there exists $L > 0$ such that

$$\|Tx - Ty\| \leq L\|x - y\| \text{ for all } x, y \in D(T).$$

T is said to be *nonexpansive* if $L = 1$ in the above inequality. T is called *λ -strictly pseudocontractive* if there exists $\lambda \in (0, 1)$ and $j(x - y) \in J(x - y)$ such that

$$\langle Tx - Ty, j(x - y) \rangle \leq \|x - y\|^2 - \lambda \|x - y - (Tx - Ty)\|^2 \text{ for all } x, y \in D(T). \quad (1)$$

The Corresponding author email: wanghj@htu.cn(Wang), songy-
isheng123@yahoo.com.cn(Song).

This work is supported by the National Natural Science Foundation of P.R. China(11071279,11171094,11271112).

T is called *pseudocontractive* if $\lambda \equiv 0$ in (1). Obviously, each λ -strictly pseudocontractive mapping is a Lipschitzian and pseudocontractive mapping with $L = \frac{\lambda+1}{\lambda}$. In particular, a nonexpansive mapping is λ -strictly pseudocontractive mapping in a Hilbert space, but the conversion may be false.

For finding a fixed point of λ -strictly pseudocontractive mapping T , a strong convergence theorem was obtained by Zhou [12] in a 2-uniformly smooth Banach space.

Theorem Z. (Zhou [12, Theorem 2.3]) Let C be a closed convex subset of a real 2-uniformly smooth Banach space E and let $T : C \rightarrow C$ be a λ -strict pseudocontraction with $F(T) \neq \emptyset$. Given $u, x_0 \in C$, a sequence $\{x_n\}$ is generated by

$$x_{n+1} = \beta_n u + \gamma_n x_n + (1 - \beta_n - \gamma_n)[\alpha_n T x_n + (1 - \alpha_n)x_n], \quad (2)$$

where $\{\alpha_n\}$, $\{\beta_n\}$ and $\{\gamma_n\}$ in $(0, 1)$ satisfy:

- (i) $\alpha_n \in [a, \mu]$, $\mu = \min\{1, \frac{\lambda}{K^2}\}$ for some constant $a \in (0, \mu)$;
- (ii) $\lim_{n \rightarrow \infty} \beta_n = 0$ and $\sum_{n=1}^{\infty} \beta_n = \infty$;
- (iii) $\lim_{n \rightarrow \infty} |\alpha_{n+1} - \alpha_n| = 0$;
- (iv) $0 < \liminf_{n \rightarrow \infty} \gamma_n \leq \limsup_{n \rightarrow \infty} \gamma_n < 1$.

Then the sequence $\{x_n\}$ converges strongly to a fixed point z of T .

Recently, Zhang and Su [13] extended Zhou's results to q -uniformly smooth Banach space. We also note that the above results excluded $\gamma_n \equiv 0$ and $\gamma_n = \frac{1}{n+1}$. Very recently, Chai and Song [1] studied the strong convergence of the modified Mann's iteration (2) with $\gamma_n \equiv 0$.

Theorem CS. (Chai and Song [1, Theorem 3.1]) Let C be a closed convex subset of a real 2-uniformly smooth Banach space E and let $T : C \rightarrow C$ be a λ -strict pseudocontraction with $F(T) \neq \emptyset$. Given $u, x_0 \in C$, a sequence $\{x_n\}$ is generated by

$$x_{n+1} = \beta_n u + (1 - \beta_n)[\alpha_n T x_n + (1 - \alpha_n)x_n], \quad (3)$$

where $\{\alpha_n\}$ and $\{\beta_n\}$ in $(0, 1)$ satisfy the following control conditions:

- (i) $\alpha_n \in [a, \mu]$, $\mu = \min\{1, \frac{\lambda}{K^2}\}$ for some constant $a \in (0, \mu)$;
- (ii) $\sum_{n=1}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty$;
- (iii) $\lim_{n \rightarrow \infty} \beta_n = 0$, $\sum_{n=1}^{\infty} \beta_n = \infty$ and $\sum_{n=1}^{\infty} |\beta_{n+1} - \beta_n| < \infty$.

Then, the sequence $\{x_n\}$ converges strongly to a fixed point z of T .

In this paper, we will deal with strong convergence of the modified Mann's iteration (2) under more relaxed conditions on the sequences $\{\alpha_n\}$, $\{\beta_n\}$ and $\{\gamma_n\}$ in $(0, 1)$,

- (i) $\alpha_n \in [0, \mu]$, $\mu = \min\{1, \frac{\lambda}{K^2}\}$ with $\liminf_{n \rightarrow \infty} \alpha_n(\lambda - K^2 \alpha_n) > 0$;
- (ii) $\lim_{n \rightarrow \infty} \beta_n = 0$ and $\sum_{n=1}^{\infty} \beta_n = \infty$;
- (iii) $\limsup_{n \rightarrow \infty} \gamma_n < 1$.

Our results obviously develop and complement the corresponding ones of Zhou [12], Song and Chai [9], Chai and Song [1], Zhang and Su [13] and others. Moreover,

our conditions are simpler, which contain $\gamma_n \equiv 0$ and $\gamma_n = \frac{1}{n+1}$ as special cases. Our conclusions may be regarded as a unification of the some existing results.

For achieving our purposes, the following facts and results are needed. Let $\rho_E : [0, \infty) \rightarrow [0, \infty)$ be the modulus of smoothness of E defined by

$$\rho_E(t) = \sup \left\{ \frac{1}{2}(\|x + y\| + \|x - y\|) - 1 : x \in S(E), \|y\| \leq t \right\}.$$

Let $q > 1$. A Banach space E is said to be q -uniformly smooth if there exists a fixed constant $c > 0$ such that $\rho_E(t) \leq ct^q$ and uniformly smooth if $\lim_{t \rightarrow 0} \frac{\rho_E(t)}{t} = 0$. Clearly, a q -uniformly smooth space must be uniformly smooth. Typical example of uniformly smooth Banach spaces is L_p ($p > 1$). More precisely, L_p is $\min\{p, 2\}$ -uniformly smooth for every $p > 1$.

Lemma 1. (Zhou [12, Lemma 1.2]) *Let C be a nonempty subset of a real 2-uniformly smooth Banach space E with the best smooth constant K , and let $T : C \rightarrow C$ be a λ -strict pseudocontraction. For any $\alpha \in (0, 1)$, we define $T_\alpha = (1 - \alpha)x + \alpha Tx$. Then,*

$$\|T_\alpha x - T_\alpha y\|^2 \leq \|x - y\|^2 - 2\alpha(\lambda - K^2\alpha)\|Tx - Ty - (x - y)\|^2 \text{ for all } x, y \in C. \quad (4)$$

In particular, as $\alpha \in (0, \frac{\lambda}{K^2}]$, $T_\alpha : C \rightarrow C$ is nonexpansive such that $F(T_\alpha) = F(T)$.

Lemma 2 was shown and used by several authors. For detail proofs, see Liu [2] and Xu [10, 11]. Furthermore, a variant of Lemma 1 has already been used by Reich in [6, Theorem 1].

Lemma 2. *Let $\{a_n\}$ be a sequence of nonnegative real numbers such that*

$$a_{n+1} \leq (1 - t_n)a_n + t_n c_n, \quad \forall n \geq 0.$$

Assume that $\{t_n\} \subset [0, 1]$ and $\{c_n\} \subset (0, +\infty)$ satisfy the restrictions:

$$\sum_{n=0}^{\infty} t_n = \infty \text{ and } \limsup_{n \rightarrow \infty} c_n \leq 0.$$

Then as $n \rightarrow \infty$, $\{a_n\}$ converges to zero.

Morales and Jung [3], in 2000, proved the following behavior for pseudocontractive mappings. Also see Song and Chen [7, 8] for more details. The same result of nonexpansive mapping was shown by Reich [5] in 1980.

Lemma 3. ([3, 7, 8]) *Let C be a nonempty, closed and convex subset of a uniformly smooth Banach space E , and let $T : C \rightarrow C$ be a continuous pseudocontractive mapping with $F(T) \neq \emptyset$. Suppose that for $t \in (0, 1)$ and $u \in C$, x_t defined by*

$$x_t = tu + (1 - t)Tx_t. \quad (5)$$

Then, as $t \rightarrow 0$, x_t converges strongly to a fixed point of T .

This following results play a key role in proving our main results, which was proved by Song and Chen [7].

Lemma 4. (Song and Chen [7, Theorem 2.3]) *Let C be a nonempty, closed and convex subset of a uniformly smooth Banach space E , and let $T : C \rightarrow C$ be a continuous pseudocontractive mapping with a fixed point. Assume that there exists a bounded sequence $\{x_n\}$ such that $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$ and $z = \lim_{t \rightarrow 0} z_t$ exists, where $\{z_t\}$ is defined by (5). Then*

$$\limsup_{n \rightarrow \infty} \langle u - z, J(x_n - z) \rangle \leq 0.$$

We also need the following results that showed by Mainge in 2008.

Lemma 5. (Mainge [4, Lemma 3.1]) *Let $\{\Gamma_n\}$ be a sequence of real numbers that does not decrease at infinity, in the sense that there exists a subsequence $\{\Gamma_{n_k}\}$ of $\{\Gamma_n\}$ such that*

$$\Gamma_{n_k} < \Gamma_{n_k+1} \text{ for all } k \geq 0.$$

Also consider the sequence of integers $\{\tau(n)\}_{n \geq n_0}$ defined by

$$\tau(n) = \max\{k \leq n; \Gamma_k < \Gamma_{k+1}\}.$$

Then $\tau(n)$ is a nondecreasing sequence verifying

$$\lim_{n \rightarrow \infty} \tau(n) = +\infty,$$

and, for all $n \geq n_0$, the following two estimates hold:

$$\Gamma_{\tau(n)} \leq \Gamma_{\tau(n)+1} \text{ and } \Gamma_n \leq \Gamma_{\tau(n)+1}.$$

Next we will show our main results.

Theorem 6. *Let C be a closed convex subset of a real 2-uniformly smooth Banach space E and let $T : C \rightarrow C$ be a λ -strict pseudo-contraction with $F(T) \neq \emptyset$. Given $u, x_0 \in C$, a sequence $\{x_n\}$ is generated by the modified Mann's iteration (2), where $\{\alpha_n\}$, $\{\beta_n\}$ and $\{\gamma_n\}$ in $(0, 1)$ satisfy:*

(i) $\alpha_n \in [0, \mu]$, $\mu = \min\{1, \frac{\lambda}{K^2}\}$ with $\liminf_{n \rightarrow \infty} \alpha_n(\lambda - K^2\alpha_n) > 0$;

(ii) $\lim_{n \rightarrow \infty} \beta_n = 0$ and $\sum_{n=1}^{\infty} \beta_n = \infty$;

(iii) $\limsup_{n \rightarrow \infty} \gamma_n < 1$.

Then the sequence $\{x_n\}$ converges strongly to a fixed point z of T .

Proof. Let $y_n = T_{\alpha_n}x_n = \alpha_nTx_n + (1-\alpha_n)x_n$. Then for each n , T_{α_n} is nonexpansive and $F(T) = F(T_{\alpha_n})$ by Lemma 1. So, the sequence $\{x_n\}$ is bounded since for given $p \in F(T) = F(T_{\alpha_n})$,

$$\begin{aligned} \|x_{n+1} - p\| &= \|\beta_n(u - p) + \gamma_n(x_n - p) + (1 - \beta_n - \gamma_n)(T_{\alpha_n}x_n - p)\| \\ &\leq \beta_n\|u - p\| + \gamma_n\|x_n - p\| + (1 - \beta_n - \gamma_n)\|T_{\alpha_n}x_n - T_{\alpha_n}p\| \\ &\leq \beta_n\|u - p\| + \gamma_n\|x_n - p\| + (1 - \beta_n - \gamma_n)\|x_n - p\| \\ &\leq \beta_n\|u - p\| + (1 - \beta_n)\|x_n - p\| \\ &\leq \max\{\|x_n - p\|, \|u - p\|\} \\ &\vdots \\ &\leq \max\{\|x_0 - p\|, \|u - p\|\}. \end{aligned}$$

Now we show $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$. It follows from Lemma 1 that

$$\|y_n - p\| = \|T_{\alpha_n}x_n - p\|^2 \leq \|x_n - p\|^2 - 2\alpha_n(\lambda - K^2\alpha_n)\|x_n - Tx_n\|^2. \quad (6)$$

Furthermore, we also have

$$\begin{aligned}
\|x_{n+1} - p\|^2 &= \|\beta_n(u - p) + \gamma_n(x_n - p) + (1 - \beta_n - \gamma_n)(y_n - p)\|^2 \\
&\leq \beta_n\|u - p\|^2 + \gamma_n\|x_n - p\|^2 \\
&\quad + (1 - \beta_n - \gamma_n)(\|x_n - p\|^2 - 2\alpha_n(\lambda - K^2\alpha_n)\|x_n - Tx_n\|^2) \\
&\leq \beta_n\|u - p\|^2 + (1 - \beta_n)\|x_n - p\|^2 \\
&\quad - 2\alpha_n(1 - \beta_n - \gamma_n)(\lambda - K^2\alpha_n)\|x_n - Tx_n\|^2 \\
&\leq \|x_n - p\|^2 - (2\alpha_n(1 - \beta_n - \gamma_n)(\lambda - K^2\alpha_n)\|x_n - Tx_n\|^2 - \beta_n\|u - p\|^2).
\end{aligned}$$

Then we obtain

$$2\alpha_n(1 - \beta_n - \gamma_n)(\lambda - K^2\alpha_n)\|x_n - Tx_n\|^2 \leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2 + \beta_n\|u - p\|^2.$$

It follows from Lemma 3 that there exist $z \in F(T)$ and $x_t = tu + (1 - t)Tx_t$ such that $\lim_{t \rightarrow 0} x_t = z$. Then we also have

$$2\alpha_n(1 - \beta_n - \gamma_n)(\lambda - K^2\alpha_n)\|x_n - Tx_n\|^2 \leq \|x_n - z\|^2 - \|x_{n+1} - z\|^2 + \beta_n\|u - z\|^2. \quad (7)$$

Following the proof technique in Mainge [4, Lemma 3.2, Theorem 3.1], the proof may be divided two cases.

Case 1. If there exists N_0 such that the sequence $\{\|x_n - z\|^2\}$ is nonincreasing for $n \geq N_0$, then the limit $\lim_{n \rightarrow \infty} \|x_n - z\|^2$ exists, and hence $\lim_{n \rightarrow \infty} (\|x_n - z\|^2 - \|x_{n+1} - z\|^2) = 0$. So by the condition (ii) and the inequality (7), it is obvious that

$$\limsup_{n \rightarrow \infty} \alpha_n(1 - \beta_n - \gamma_n)(\lambda - K^2\alpha_n)\|x_n - Tx_n\|^2 = 0.$$

It follows from the conditions (i), (ii) and (iii) that

$$\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0. \quad (8)$$

Then by Lemma 4, we obtain

$$\limsup_{n \rightarrow \infty} \langle u - z, J(x_{n+1} - z) \rangle \leq 0. \quad (9)$$

Finally, we show that $x_n \rightarrow z$. Indeed, since

$$\begin{aligned}
\|x_{n+1} - z\|^2 &= \langle (\beta_n u + \gamma_n x_n + (1 - \beta_n - \gamma_n)Tx_n) - z, J(x_{n+1} - z) \rangle \\
&\leq \beta_n \langle u - z, J(x_{n+1} - z) \rangle + \gamma_n \|x_n - z\| \|J(x_{n+1} - z)\| \\
&\quad + (1 - \beta_n - \gamma_n) \|Tx_n - z\| \|J(x_{n+1} - z)\| \\
&\leq \beta_n \langle u - z, J(x_{n+1} - z) \rangle + (1 - \beta_n) \|x_n - z\| \|x_{n+1} - z\| \\
&\leq \beta_n \langle u - z, J(x_{n+1} - z) \rangle + (1 - \beta_n) \frac{\|x_n - z\|^2 + \|x_{n+1} - z\|^2}{2},
\end{aligned}$$

then, we have

$$\|x_{n+1} - z\|^2 \leq (1 - \beta_n) \|x_n - z\|^2 + 2\beta_n \langle u - z, J(x_{n+1} - z) \rangle. \quad (10)$$

So, an application of Lemma 2 onto (10) yields that $\lim_{n \rightarrow \infty} \|x_n - z\| = 0$.

Case 2. Assume that there exists a subsequence $\{\|x_{n_k} - z\|^2\}$ of $\{\|x_n - z\|^2\}$ such that $\|x_{n_k} - z\|^2 < \|x_{n_k+1} - z\|^2$ for all $k \geq 0$. Let

$$\Gamma_n = \|x_n - z\|^2 \text{ and } \tau(n) = \max\{k \leq n; \Gamma_k < \Gamma_{k+1}\}.$$

It follows from Lemma 5 that $\tau(n)$ is a nondecreasing sequence verifying

$$\lim_{n \rightarrow \infty} \tau(n) = +\infty$$

and for n large enough,

$$\Gamma_{\tau(n)} \leq \Gamma_{\tau(n)+1}, \quad \Gamma_n = \|x_n - z\|^2 \leq \Gamma_{\tau(n)+1}. \quad (11)$$

In light of Eq. (7), we have

$$2\alpha_{\tau(n)}(1 - \beta_{\tau(n)} - \gamma_{\tau(n)})(\lambda - K^2\alpha_{\tau(n)})\|x_{\tau(n)} - Tx_{\tau(n)}\|^2 \leq \beta_{\tau(n)}\|u - z\|^2,$$

and so by the condition (i),(ii) and (iii), we have

$$\lim_{n \rightarrow \infty} \|x_{\tau(n)} - Tx_{\tau(n)}\| = 0.$$

Then as $n \rightarrow \infty$,

$$\begin{aligned} \|x_{\tau(n)+1} - Tx_{\tau(n)}\| &\leq \beta_{\tau(n)}\|u - Tx_{\tau(n)}\| + \gamma_{\tau(n)}\|x_{\tau(n)} - Tx_{\tau(n)}\| \\ &\quad + (1 - \beta_{\tau(n)} - \gamma_{\tau(n)})(1 - \alpha_{\tau(n)})\|x_{\tau(n)} - Tx_{\tau(n)}\| \rightarrow 0. \end{aligned}$$

Since

$$\begin{aligned} \|x_{\tau(n)+1} - Tx_{\tau(n)+1}\| &\leq \|x_{\tau(n)+1} - Tx_{\tau(n)}\| + \|Tx_{\tau(n)} - Tx_{\tau(n)+1}\| \\ &\leq \|x_{\tau(n)+1} - Tx_{\tau(n)}\| + \|x_{\tau(n)} - x_{\tau(n)+1}\| \\ &\leq 2\|x_{\tau(n)+1} - Tx_{\tau(n)}\| + \|x_{\tau(n)} - Tx_{\tau(n)}\|, \end{aligned}$$

we have

$$\lim_{n \rightarrow \infty} \|x_{\tau(n)+1} - Tx_{\tau(n)+1}\| = 0. \quad (12)$$

Then by Lemma 4, we obtain

$$\limsup_{n \rightarrow \infty} \langle u - z, J(x_{\tau(n)+1} - z) \rangle \leq 0. \quad (13)$$

Using the similar proof techniques of Case 1, the only modification is that n is replaced by $\tau(n)$, we have

$$\|x_{\tau(n)+1} - z\|^2 \leq (1 - \beta_{\tau(n)})\|x_{\tau(n)} - z\|^2 + 2\beta_{\tau(n)}\langle u - z, J(x_{\tau(n)+1} - z) \rangle. \quad (14)$$

Together with (11), we have

$$\Gamma_{\tau(n)} \leq \Gamma_{\tau(n)+1} \leq (1 - \beta_{\tau(n)})\Gamma_{\tau(n)} + 2\beta_{\tau(n)}\langle u - z, J(x_{\tau(n)+1} - z) \rangle,$$

and so,

$$\Gamma_{\tau(n)} = \|x_{\tau(n)} - z\|^2 \leq 2\langle u - z, J(x_{\tau(n)+1} - z) \rangle.$$

Along with (13), we have

$$\lim_{n \rightarrow \infty} \Gamma_{\tau(n)} = \lim_{n \rightarrow \infty} \|x_{\tau(n)} - z\|^2 = 0.$$

It follows from (14), (13) and the condition (ii) that

$$\lim_{n \rightarrow \infty} \Gamma_{\tau(n)+1} = \lim_{n \rightarrow \infty} \|x_{\tau(n)+1} - z\|^2 = 0.$$

Now it follows from (11) that

$$\lim_{n \rightarrow \infty} \Gamma_n = \lim_{n \rightarrow \infty} \|x_n - z\|^2 = 0.$$

The proof is completed. \square

Clearly, Theorem 6 contains $\gamma_n \equiv 0$ and $\gamma_n = \frac{1}{n+1}$ as special cases. So the following result is obtained easily.

Corollary 7. Let C be a closed convex subset of a real 2-uniformly smooth Banach space E and let $T : C \rightarrow C$ be a λ -strict pseudo-contraction with $F(T) \neq \emptyset$. Given $u, x_0 \in C$, a sequence $\{x_n\}$ is generated by the modified Mann's iteration (3), where $\{\alpha_n\}$ and $\{\beta_n\}$ in $(0, 1)$ satisfy:

- (i) $\alpha_n \in [0, \mu]$, $\mu = \min\{1, \frac{\lambda}{K^2}\}$ with $\liminf_{n \rightarrow \infty} \alpha_n(\lambda - K^2\alpha_n) > 0$;
- (ii) $\lim_{n \rightarrow \infty} \beta_n = 0$ and $\sum_{n=1}^{\infty} \beta_n = \infty$.

Then the sequence $\{x_n\}$ converges strongly to a fixed point z of T .

Using the same proof techniques as Theorem 6, we easily obtain the following result. Since the only difference is that $\alpha_n(\lambda - K^2\alpha_n)$ is replaced by $\alpha_n(q\lambda - C_q\alpha_n^{q-1})$ in its proof, so we omit its proof.

Theorem 8. Let C be a closed convex subset of a real q -uniformly smooth Banach space E ($q > 1$) and let $T : C \rightarrow C$ be a λ -strict pseudo-contraction with $F(T) \neq \emptyset$. Given $u, x_0 \in C$, a sequence $\{x_n\}$ is generated by the modified Mann's iteration (2) or (3), where $\{\alpha_n\}$, $\{\beta_n\}$ and $\{\gamma_n\}$ in $(0, 1)$ satisfy:

- (i) $\alpha_n \in [0, \mu]$, $\mu = \min\{1, \{\frac{q\lambda}{C_q}\}^{\frac{1}{q-1}}\}$ with $\liminf_{n \rightarrow \infty} \alpha_n(q\lambda - C_q\alpha_n^{q-1}) > 0$;
- (ii) $\lim_{n \rightarrow \infty} \beta_n = 0$ and $\sum_{n=1}^{\infty} \beta_n = \infty$;
- (iii) $\limsup_{n \rightarrow \infty} \gamma_n < 1$.

Then the sequence $\{x_n\}$ converges strongly to a fixed point z of T .

REFERENCES

1. X. Chai and Y. Song, *Convergence theorem for an iterative algorithm of lambda-strict pseudocontraction*, Fixed Point Theory and Applications 2011, 2011:95 (6 December 2011).
2. L.S. Liu, *Ishikawa and Mann iteration process with errors for nonlinear strongly accretive mappings in Banach spaces*, J. Math. Anal. Appl. 194(1995) 114–125.
3. C.H. Morales, J.S. Jung, *Convergence of paths for pseudo-contractive mappings in Banach spaces*, Proc. Amer. Math. Soc. 128(2000), 3411–3419.
4. Paul-Emile Mainge, *Strong Convergence of Projected Subgradient Methods for Nonsmooth and Nonstrictly Convex Minimization*, Set-Valued Anal 16(2008) 899–912.
5. S. Reich, *Strong convergence theorems for resolvents of accretive operators in Banach spaces*, J. Math. Anal. Appl. 75(1980) 287–292.
6. S. Reich, *Constructive techniques for accretive and monotone operators*, in: Applied Nonlinear Analysis (Academic Press, New York, 1979), pp. 335–345.
7. Y. Song, R. Chen, *Convergence theorems of iterative algorithms for continuous pseudocontractive mappings*, Nonlinear Anal. 67(2007), 486–497.
8. Y. Song, R. Chen, *An approximation method for continuous pseudocontractive mappings*, J. Inequal. Appl. 2006(2006), Article ID 28950, 9 pages. doi:10.1155/JIA/2006/28950.
9. Y. Song, X. Chai, *Halpern iteration for firmly type nonexpansive mappings*, Nonlinear Anal. 71(10)(2009) 4500–4506.
10. H.K. Xu, *An iterative approach to quadratic optimization*, J. Optim. Theory Appl., 116(2003) 659–678.
11. H.K. Xu, *Iterative algorithms for nonlinear operators*, J. London Math. Soc. 66 (2002) 240–256.
12. H. Zhou, *Convergence theorems for λ -strict pseudo-contractions in 2-uniformly smooth Banach spaces*, Nonlinear Anal. 69(9)(2008), 3160–3173.
13. H. Zhang and Y. Su, *Convergence theorems for strict pseudo-contractions in q -uniformly smooth Banach spaces*, Nonlinear Anal. 71(2009), 4572–4580.